

(12)

Notodelphyoidea - Buproridae.

Buprorus, valid -

Thorell, 1859.

B. lovénii ^(M).

Syn. Buprorus Thorell, 1859, p. 61.

Buprorus Sars, 1921, p. 61.

Βούρρωπος = having the face of an ox.

Body (♀) sub-saccate, oblong, & head small = rounded posteriorly: anal laminae locking. No eyes.

1st. antennae short, 3-jointed: 2nd pair set with stout, curved setae: Mandibles narrow, with sharp teeth on & inner margin = no palps.

Maxillae compressed; 1st & 2nd pairs composed of a major basal part & 2 smaller joints, sub-parallel 3rd. pair of a single joint, armed with obtuse, parallel setae. Legs short & wide, first 4

pairs biramous; rami with spines or setae at & apex = endopod simple, exopod 2-jointed.

5th pair rudimentary, simple, sub-conical.

(Thorell, 1859, p. 61)

In respiratory sac of Ascidian

Notodelphyoidea - Buproridae.
Buprorus, valid -

Thorell, 1859.

B. lovénii ^(M)

Syn. Buprorus Thorell, 1859, p. 61.

Buprorus Sars, 1921, p. 61.

Βούπρωπος = having the face of an ox.

Body (♀) sub-saccate, oblong, & head small - rounded posteriorly; anal laminae locking. No eyes.
1st. antennae short, 3-jointed: 2nd pair set with stout, curved setae: Mandibles narrow, with sharp teeth on & inner margin = no palps.
Maxillae compressed; 1st & 2nd pairs composed of a major basal part & 2 smaller joints, sub-parallel; 3rd. pair of a single joint, armed with obtuse, parallel setae. Legs short & wide, first 4 pairs biramous; rami with spines or setae at & apex = endopod simple, exopod 2-jointed.
5th pair rudimentary, simple, sub-conical.

(Thorell, 1859, p. 61)

In respiratory sac of Ascidians.

BUPRORUS THORELL, 1859

2

- ILLG + DUDLEY 1980: 6, 9-14, 31-32, KEY TO SPP.; 155
- GOTTO 1993: 51, DIAGNOSIS

BUPRORUS CAUDATUS ILLG + DUDLEY 1980

--, ILLG + DUDLEY 1980: 32-35, FIG. 7, N. SP. ♀ -

HOST STYELA CORIACEA, OFF SOUTHERN CALIFORNIA

-- OISHI 1991: 52 -

Buproridae - Notodelphyoidea

Buprorus loveni Thorell, ♀.

- —, Thorell, 1859, p. 63: pl. 10, fig. 14.
— —, Aurivillius, 1883, p. 33: pl. 1, f. 13.
— —, Nordenskjöld, 1885, p. 281.
— —, Thorell, 1859, a, p. 340.

Host, Ascidia obliqua: A. mentula:
A. aspersa, Norway: Bohuslän:
Faroes: Mediterranean

— — Aurivillius, 1882, no. 3, p. 63; pl. 5, f. 13.

— — Wilson, 1920, p. 14 L

— — Sars, 1921, p. 62: pl. 30

— — Hausen, 1923, p. 23.

— — Stephensen, 1929, p. 7.

-- --, Bresciani et Lutzen 1962. p. 373

-- ILLG & DUDLEY 1980: 32, IN KEY; GIVES LIST OF HOSTS; 39

-- DUDLEY + ILLG 1991: 11-, FIGS. 5A, 7; REVIEW RECORDS

MASS., N. H., MAINE

-- OCISHI 1991: 62

-- GOTTO 1993: 51-, FIG. 9M

Buproridae Notodelphyoidea

Buprorus nordgaardii G.O. Sars,

— — Sars, 1921, p. 90 nomen nudum.

— —, ILLG + DUDLEY 1980: 11, 32, IN KEY; GIVES HOST LISTS; 35

— — OOISHI 1991: 62

BRUPRORUS PRAZINOIDES

-- OF SEWELL, 1949 = ANOMOPSYLLUS -, ILLG + DUDLEY 1980: