

in

Lernaeopodoida - Lernaeopodidae.
Siphonostomatoida

Clavellodes valid

Wilson, 1915 —

C. rugosa [Ⓛ].

Syn. Clavellodes Wilson, 1915, p. 689.
Anchorella (pars) various authors.

Clavella and ending εἶδος = likeness.

Clavellodes Wilson, 1915

- 2 -

- , Kabata, 1967, p. 1298
- CASTRO + BAEZA 1984: ~~68~~ 3-4
- KABATA 1988: 42, 68; ONLY SPECIES IS C. RUGOSUS
- KABATA 1988b: 7
- KABATA 1992: 132, DIAGNOSIS

Clavellodes canthari (Heller, 1865) - see Alella canthari

Lernaeopodidae - Lernaeopodoida

Clavellodes ditrematis Yamaguti ♂♀.

— — Yamaguti, 1939, p. 564: pl. 54, figs. 207-214.

--, Kabata, 1969, p. 323

Host. Gills, Hexagrammos otakii Jordan & Starks.

Tarumi, Japan.

Clavellodes intermedia (Quider). ♂. ♀.

Anchorella —, Quider, 1906, p. 29: pl. 3, figs. 37-44

Clavellodes —, Wilson, 1915, pp. 669, 690.

— — Wilson, 1920, p. 72.

—, Kabata, 1969, p. 322.

Host. Mouth cavity of a Nothostemna.

Antarctic.

Clavellodes macrotrachelus (Brian) D.F.
1906

Anchorella uncinata, Brian, 1908, p. 221.

— couthari, Brian, 1903a, p. 82

— —, —, 1903b., p. 9. [pl. 21, figs. 1-4

Clavella macrotrachelus, —, 1906, p. 116: pl. 8, fig. 5: ^

Host Gills of Sargus vulgaris + S. roudoletii.

Mediterranean.

Clavellodes m—, Wilson, 1915, pp. 669 & 690.

— — Brian, 1924, p. 52.

— —, Nunes-Ruivo, 1956, p. 33. On Smaris melancurus
Baía Farta, Benguela, Angola.

--, Kabata, 1969, p. 322.

--, CAILLET & RAIBAUT, 1979: 993-996
EX DIPLODUS SARGUS (FISH)

Clavellodes macrotrachelus

card 2

- , Kawatow, et al., 1980, p. 199
- CAILLET + RAIBAUT 1979: 223-226, FIG. 1, HOST
DIPLODUS SARGUS
- COSTE ET AL. 1982: 259

Clavellodes pagelli (Krøyer, 1863)

--, Kabata, 1969, p. 323.

--, Kabata, 1972, p. 216

-- KENSLEY & GRINDLEY 1973: 127

Clavellodes pterobrachiata Kabata, 1968, pp. 505,

n. sp., Kabata, 1968, pp. 505, 518-9, fig 1 D-H.

--, Kabata, 1969, p. 323

Clavellodes rugosa (Krøyer) ♂, ♀.

[fig. 14a-c.]

Anchorella —, Krøyer, 1837, p. 284, pl. II, fig. 7; pl. III,

— —, Milne Edwards, 1840, p. 519.

— —, Baird, 1850, p. 338: pl. 35, fig. 8.

— —, White, 1850, p. 130 - 1857, p. 331.

Host. Gills of cat-fish Auarrhichas lupus.
also on Radus cellarius (Baird)Greenland: Iceland: Faroes: British Isles:
New England: Belgium: Arctic Ocean.

— — . P. J. van Beneden, 1851, p. 114: pl. 6, figs. 7-10.

— — . ————, 1861a, p. 152.

— — . Krøyer, 1863, p. 383.

— — . Olsson, 1869, p. 44: B-Smith, 1899, p. 503.

— — . Vogt — 1877, b, pl. 10, fig. 9.

— — . T. Scott, 1900, p. 176: pl. 8, figs. 45-48.

— — . Stephensen, 1913, p. 338 figs. 190-198

Clavellodes —, Wilson, 1915, p. 690, pl. 27, fig. 6: pl. 51

Anchorella — Norman & Brady, 1909, p. 158,

— — Bainbridge, 1909, p. 55: pl. 10, figs. 28-32: pl. 11, fig. 33

— — T. Scott, 1906, b, p. 374

— — Wilson, 1920, p. 7L

— — Wilson, 1932, p. 516: fig. 305.

Clavellodes rugosus (Krøyer, 1837) card 2

--, Kabata, 1969, pp. 316-318, figs. 24-35.

see Anchorella rugosa (Kroyer, 1837 & Clavella rugosa

--, Polianskii, 1955, p. 94-

(Kroyer) T. Scott, 1900

--, Stephensen, 1940, p. 13, ~~якхуякхуя~~ Clavella r. is a syn

--, Ho, 1977, p. 8, 12, fig. 9

--, Kabata & Cousens, 1972, passim

--, AULOQUE, 1899: 206 - as ANCHORELLA RUGOSA.

-- KABATA 1988: 68, FIG. 33A-E; HOST LIST.

-- KABATA 1992: 132; FIG. 29F-J, SYNONYMY, HOST LIST