

Galoiopotes, valid,
Stp. & Ltk., 1861,

Caligoidea
Eurypheridae Caligidae.
G. hygomianus.

A species was described in 1889
by Thomson under the name
Twepeophtheirus huttoni.

Γλοιός = sticky, Knarish : πότης = drinker, or
tippler.

Carapace large oval, shield shaped -
anterior thorax seg'ts joined with the head -
Frontal plates distinct, without lunules -
Fourth thorax seg't free, with 2 dorsal plates
partly covering v gen. seg't. The latter large,
and produced backward on either side of the
abdomen in form of an elongated, curved
process, having styliform appendages on the
outer border, + serrated at v edge.
Abdomen long, 2 jointed. Anal laminae lance-
shaped - First + fourth legs uniramous, 2nd +
3rd biramous. Egg-cases + eggs as in Caligus.

GLOIOPOTES

2

- , CRESSEY, 1961: 1-2, KEY TO FEMALES p. 2;
9-13 DISCUSSION
- DOJIRI, 1983: 192-195
- CRESSEY ET AL. 1983: 232-
- DOJIRI & CRESSEY 1991: 605-

GLOIOPOTES AMERICANUS CRESSEY, 1967

-- CRESSEY, 1967: 2-5 ♀♂, 6, 7, 8, 10, 13; FIGS. 1-24; Pl. 1, 2

EX ISTIOPHORUS AMERICANUS

--, CRESSEY + LACHNER, 1970: 314

-- CRESSEY ET AL. 1983: 240-, HOSTS

Gloiopotes auricularius Barnard

1957-- Barnard, K.H., Ann. Mag. Nat. Hist. Ser. 12,
vol. X, part 1, p. 11-12, fig. 8.

~~Leois, 1967: 66~~

GLOIOPOTES ADRICULATUS BARNARD

--, LEWIS, 1967: 66

Gloiopotes costatus, New Species, ♂♀

— — Wilson, 1919, p. 313; pl. 21, f. 1-9.

--, Pillai, 1964, p. 63

--, Lewis, 1967: 65

Host, Outside surface of "Swordfish"
Catalina Island, California.

Carigidae - Carigoida

Gloiopotes crassus Wilson & Bere.

— — Bere, 1936,

Host, On Rhombochirus osteochir and Tetrapturus
imperator

Florida.

Gloiopotes Luttoni, Thomson, Caligidae ♂.♀. ①

Lepeophtheirus Luttoni, Thomson, 1889, p. 354.
pl. XXVIII, fig. 10a-c; pl. XXIX, fig. 1a-m.

Gloiopotes Luttoni, B. Smith, 1899, p. 458.

-- --, Hewitt 1964, pp. 85-96, fig. 1-16

Host. Nistiophorus Herscheli, N. Zealand
Nistiophorus, spp. Madras.

Gloiopotes huttoni

(2)

-- -- Cressey, 1967, p. 6, 2, 4, 6-7, 8, 10, 13; FIG. 37-37;

-- -- Kabata, 1966, p. 568.

Pl. 3, 4

-- --, Cressey + Lachner, 1970, p. 314, 317

--, Lewis, 1967, pp. 57, 58, 60, 62, 63, 65, ^{100;} figs. 22-24

--, Hewitt & Hine, 1972, p. 89

-- CRESSEY ET AL. 1983: 240-, HOSTS

-- Ho + I. H. KIM 1996: 286-289, SEA OF JAPAN;
HOST XIPHIAS GLADIUS. SYN. G. LONGICAUDATUS.

Gloiopotes lygomianus, Stp. & Ltk. ♀. Euryphoridae
Saligidae.

— —, Stp. & Ltk, 1861, p. 363; pl. V. fig. 9.

— —, B-Smilk, 1899, p. 458. (in list)

— —, Stebbing, 1901, p. 670; pl. LXXIV, fig. A.
(descr. + fig. of ♂).

— — Cressy, 1967, p. 8

Host, An unknown fish.

Rubiana, New Georgia,

Atlantic. (Stp. + Ltk.)

from tail of albacore - Stebbing.

-- -- Shiino, 1960. pp. 533-539, fig. 4-6

--, Lewis, 1969, fig. 2b

--, Lewis, 1967, pp. ~~59, 58, 60, 62, 63, 65, figs. 21-24~~
pp. 66, 67, 69, 71, 72, figs. 25-27

--, Kazachenko, 1975, p. 215

--, CRESSEY, 1967: 2, 8-9 ♀♀, 10, 13; FIG. 40-42

--, CRESSEY + CRESSEY 1980: 38; Figs. 79c-d.

- Dojiri, 1983: 195-204, fig. 71-77.

Gloiopotes hygomianus Stp. & Ltk. 1861

card 2

-- Cressey et al. 1983: 228-, fig. 1f; HOSTS

-- ZWERNER + LAWLER 1972: 84, CHESAPEAKE BAY,
HOST ACANTHOCYBIUM SOLANDER!

Gloiopotes longicaudatus (Marukawa) 1925

= Caligus longicaudatus

--- --- Shiino 1963. pp. 343-344.

-- --, Heegaard 1962. p. 174, fig. 151-153

-- --, Ho 1963. pp. 87-90, fig. 6-10

-----, Shiino 1954, art 15, pp. 273-278, fig. 1

--, Pillai, 1964, p. 63

--, Lewis, 1967: 65, 66

---, Pillai, 197?, p. 1614-1615, fig. 125

Galviopotes ornatus, Wilson, ♀ ♂

Galigidae.

- —, Wilson, 1905, a, p. 127.
— —, M. J. Rathbun, 1905, p. 93.
— —, Wilson, 1907, p. 699: pl. 17, figs. 23-34.
— —, Wilson, 1932, p. 415: fig 262. ♂ (WOODS HOLE)
— —, Wilson 1936 (Mem. Soc. Cub. Hist. Nat). On Tetrapturus amplus Poey.
Host, Outside surface of Swordfish.
— —, Cressey, 1961, p 5 Atlantic.
— —, Cressey & Lachner, 1970, p. 314

--, Pillai, 1964, p. 63

--, Lewis, 1967: 65, 66

--, CRESSEY, 1967: 2, 4, 5-6 ♂♂, 7, 8, 10, 13; FIG. 25, 26; PL. 1, 2

-- CRESSEY ET AL. 1983: 240-1, HOSTS

-- ZWERNER + LAWLER 1972: 84, ^{OFF} CHESAPEAKE BAY,

HOST MAKAIIRA NIGRICANS

Caligidae Caligoida

Gloiopotes watsoni Kirtisinghe, ♂♀. ①

— — Kirtisinghe, 1934, p. 167: text-figs. 1-17.

-- Cressey, 1967, p. 7.

--, Cressey + Lachner, 1970, p. ~~315~~ 315

--, Pillai, 1964, p. 63.

Host. Skin of Histiophorus gladius (Sailfish)
Ceylon.

--, CRESSEY, 1967: 2, 4, 7-8 ♂♂, 10, 12, 13; FIGS. 38-39; PL. 3, 4

Gloiopotes watsoni Kirtisinghe 1934

card 2

-- Kensley & Grindley 1973: 92, from Makaira indica and

Makaira audax, Cape, South Africa.; p. 124.

-- CRESSEY ~~ET AL.~~ 1983: 240-, HOSTS
Collette + Russo

Caligiidae - Caligoidae
Gloiopotes sp. Yamaguti ♂♀.

— — Yamaguti, 1936, ^{part 3,} p. 4: pl. 2, f. 20; pl. 3, figs. 21-35.

Host Skin of Tetrapturus mitsukurii Jordan & Snyder
Japan.

GLOIOPOTES SP.

-- BONNET 1948: 7, HAWAII; HOST ACANTHOCYBIUM
SOLANDRI.