

Clausiidae — Cyclopoida.
POECILOSTOMA FOIDA

Hippomolgus.

G.O.Sars, 1917

H. furcifer^(M).

Syn. Hippomolgus Sars, 1917, p. 147.

c/Ιππός = horse and μολύσ = hide.

Body not very cyclopoid in shape; anterior division only slightly dilated. Caudal rami much produced. First antennae short, 6-jointed; posterior pair feeble. Mxpds. not fully developed. Swimming legs coarsely made, 1st. pair not modified, last pair 1-jointed.

HIPPOMOLGUS

SARS

(2)

- , BOQUET & STOCK 1957A: p.213, key to genera
- , HUMES 1987a: 655
- TODA ET AL 1992: 109
- HO & WARDLE 1992: 43-44
- HUMES & Huys 1992: 1378, IN KEY

Hippomolgus cognatus

-- Humes and Ho, 1967, p. 14

Hippomolgus dubia (Thompson & Scott, 1903)

see Hersiliodes dubia : . . .

Clausidiidae - Cyclopoida

Hippomedes furcifer G.O. Sars, ♀.

— — Sars, 1917, p. 148: pl. 82.

-- VERVOORT & RAMIREZ 1966: 202

Habitat. Norway Fjords.

Hippomolagus latipes

--- Humes and Ho, 1967, p. 10