

in) SABELLIPHILIDAE- POECILOSTOMA Cyclopoida.
Modiolicola, TOIDA Tictonomolgidae.
Aurivillius, 1883, M. insignis.⁽¹⁾

Syn. Modiolicola Aurivillius, 1883, p. 39.
Modiolicola G.O. Sars, 1918, p. 173.

Modiolus and colo, to inhabit.

Cephalothorax broadly ovate, 6 segts, & 2nd marked off from head by a shallow suture: abdomen narrowed toward apex, 5 segts in ♀, & 1st wider than in thorax, with convex sides -
1st antennal filiform, as long as head, 7-jointed,
2nd pair a little shorter, simple 4-jointed, with curved spines at tip. Mandibles with a broad basal part, & a slender seta at tip.

1st. mxdps 3-jointed, ending in a slender seta: 2nd pair 3-jointed, stouter, & last joint with a few spines at tip.
4 pairs biramous feet: same 3-jointed: 5th pair rud.

1-jointed.

Paired egg-sacs affixed to v sides "1st. abdom. segt
(gen. segt + part of thorax)

(Aurivillius, 1883, p. 39)

Modiolicola

- Humes & Ho, 1966, p. 10
- HUMES + STOCK 1973: 47 - 48, DIAGNOSIS, KEY TO ♀♀
- GOTTO 1993: 79, DIAGNOSIS, KEY

Modiolicola Aurivillius, 1882

- Illg & Humes, 1971, p. 577
- BOCQUET & STOCK 1959: 238

MODIOLICOLA AVDEEV I. H. KIM, 1995

-- I. H. KIM 1995: 315-321, FIGS. 1-3, N. JR., SEA OF
JAPAN OFF KOREA; EX Modiolus affinis

Modiloicola bifida Tanaka, 1961 (sic, Modiolicola) ①

n. sp. Tanaka, 1961, pp. 266-269, pl. 32, figs. 6-11;
pl. 33, figs. 1-12; pl. 34, figs. 1-9;
pl. 35, figs. 1-10

--, Ko, Murakami, & Daiku, 1962, p. 114-117

--, Ko, 1969, p. 83, 86-87, 90-91, fig. 1, tabs. 1, 3

--, Ko, 1969, p. 5

- bifidus, Do + KAJIHARA, 1984: 5, 10-15, FIGS. 21-45 -
JAPAN

-- COSTANZO 1984: 405

- BIFIDUS, HUMES + STOCK 1973: 50; syn. PARENTHESIS
sp? OF KO, MURAKAMI + DAIDU, 1962.

Modiloicola bifida

see Modolicola

--, ANDREEV, 1977a: 34-37, 40; FIG. 1-19

Modiolicola bifidus Tanaka, 1961

card 2

- Ho & Kim 1991b: 369, 370, 377-378, 395, Fig. 14A-H,
ex Tapes philippinarum, Scapharca broughtoni, and
Scapharca subcrenata, various localities, Korea
— I. H. Kim, 1995: 315, 316, 320

MODIOLICOLA GRACILICAUDUS AVDEEV, 1977

-- AVDEEV, 1977a: 37-405 FIG. 20-37; EK

CRENOMYTILUS GRAYANUS (MOLLUSC), ^{Possiet} BAY

--, HO, 1980, p. 298-300, figs. 2a-f, 3a-1, SADO ISLAND

-- DO + KAJIHARA 1984:5

-- HO & KIM 1991b: 369, 370, 378-379, 395, FIG. 14I-M,

EX MYTILUS CORUSCUS, KOREA, GANGREUNG + SOKCHO

-- I. H. KIM 1995: 320

Lichomolgidae - Cyl.,
Modiolicola gracilis Wilson, ♀. ①

— — Wilson, 1935, p. 781; pl. 25, figs. 8-12

--, Tanaka, 1961, p. 266

--, AODEEV, 1977a: 40

-- DO & KAJIHARA 1984: 5

-- AODEEV 1987: 612

Host. Gills of Mytilus edulis, California.

-- HUMES & STOCK 1973: 50-55, FIGS. 28-30, RECORDS-

CALIFORNIA; HOST MYTILUS CALIFORNIANUS, NOT
M. EDULIS.

MODIOLICOLA GRACILIS C. B. WILSON, 1935 (2)
-- CACERRS - MARTINEZ ET AL. 1996: 667-671,
BAJA CALIFORNIA, MEXICO

Lichomolgidae - Cyclopoida

Modiolicola inermis Canu.

— — Canu, 1891, p. 436: 1892, p. 240: fig. 19.

— — Sars, 1918, p. 173: pl. 97.

Lichomolgas maximus Thompson, 1893, p. 208: pl. 35, f. 1-11.

Modiolicola inermis, Canu, 1899, p. 73.

—, Audeev, 1977a: 41

Habitat. Gill cavity, Modiola & Mytilus.

-- Monod + Dollfus, 1932, p. 152.

--, Tanaka, 1961, p. 261, 266

-- AUDEREU 1967:612

Modiolicola inermis Canu, 1892

card 2

Paranthessius (M.) -, Bocquet & Stock, 1957:427

M. -, Humes & Stock 1973:55-56, fig. 31, 2 localities-

Roscoff, France; give synonymy; host Pecten opercularis

PARANTHESSIUS -, Stock 1960: 247-248, BANYULS, FRANCE,
ASSOC. PECTEN OPERCULARIS F. AUDOUINI

M. -, Gotto 1993: 80-, FIG. 16G-K

Modiolicola inflatipes - see Herrmannella

Tichonolgiidae - Cyclopoida.

Modiolicola insignis Aurivillius ♀♂.

p. 39: pl. 4, figs. 1-8 &

-- Aurivillius, 1883, p. 10: pl. 2, figs. 1-10 ♀

-- Aurivillius, 1882, no. 8, p. 43: pl. 13, figs. 1-8.

-- Aurivillius, 1882, No. 3, p. 40: pl. 6, figs. 1-10.

-- T. Scott, 1893, p. 207: pl. 4, figs. 13-24.

-- Hamond, 1973, p. 348.

Host. Gill plates of Modiola vulgaris
and Mytilus edulis.

Norway -

- — T. Scott, 1906, b, p. 354.
— — G. O. Sars, 1918, p. 173 : pl. 97.
— — Raffaele + Monticelli, 1885, p. 302 : figs. 13-16.
— — Canu, 1891, p. 482 : 1892, p. 238 : pl. 24, figs. 14-28.
— — Canu, 1899, p. 73.
— — Monod + Dollfus, 1932, p. 152.
— , Stock & Vos, 1960, p. 26x 206
— , Humes & Stock, 1972, p. 123
— , Tanaka, 1961, p. 266
— , Avdeev, 1977a : 40
— Do + KAJIHARA 1984: 5

(2)

MODIOLICOLA INSIGNIS AURIVILLIUS

-- COSTANZO 1984: 393-905, FIGS. 1-7, DEVELOPMENT;

HOST MYTILUS GALLOPROVINCIALIS

-- AUDEREU 1987: 612

-- HUMES & STOCK 1973: 48-50, FIG. 27, GIVE SYNONYMY;
BREST, FRANCE, EX MODIOLUS BARBATUS

-- IZAWA 1991: 72-

-- GOTTO 1993: 82-, FIGS. 17A-D

-- STOCK 1993: 62, EX MYTILUS EDULIS, NETHERLANDS

-- CACERES - MARTINEZ ET AL. 1996: 671

-- A. SCOTT 1896: 153, GREAT BRITAIN, EX MYTILUS MODIOLUS

Modiolicola jamaicensis Wilson - see Ascidioxynus

+ = MODIOLICOLA

Lichomolgidae - Cyclopoida.

Lichomolgas maximus I.C.Thompson, ♀.

↳ — I.C.Thompson, 1893, p. 208 : pl. 35, f. 1-11.

↳ — T.Scott, 1896, b, p. 62.

Pseudanthessius m- Thompson & Scott, 1903, p. 276 : pl. 14, f. 1-11.

HERRMANELLA MAXIMA, T. Scott 1906 : 354

Habitat. English Seas. Washed from gills
of Pecten maximus, P. opercularis

Modiolicola maxima (Thompson 1893) 2

--, Tanaka, 1961, p. 261; 266 (mixima)

--, Audeev, 1977a: 41

-- Gotto 1966: 194

- MAXIMUS, Audeev 1987: 612

- MAXIMUS, Humes & Stock 1973: 56-61, FIGS. 32-34,
4 RECORDS - FRANCE, ex PECTEN MAXIMUS & P.
OPERCULARIS F. AUDOUINI; GIVE SYNONYMY.

- - Stock 1960: 248

- MAXIMUS, Gotto 1993: 80-, FIG. 16 L-O

Modiolicola peronidius Avdeev, 1987

-- Avdeev, 1987:608, 612, figs. 2(7,8),3; n. sp. ♀ only,
host Peronidia venulosus, Peter the Great Bay, Sea of
Japan

-- I.H. Kim 1995: 320

Modiolicola trabalis Humes

1958---pp. 315-319, figs. 250-268.

--, Tanaka, 1961, p. 266

--, AVDEEV, 1977a : 40

-- AVDEEV 1987: 612

-- HUMES & STOCK 1973: 61