

Diaptomidae - Calanoida.

Pseudolovenula

Marukawa, 1921.

P. magna ^(M.).

Syn. Pseudolovenula, Marukawa, 1921, p. 13.

♀. Body slender, fusiform in metasome, apparently 6-seg'd; head separated from 1st segt. 5th segt rounded at corners. Urosome 3-seg'd. 1st antennae 24-seg'd - reaching 5-segts beyond tips of caudal rami. Exopod of 2nd antennae 8-seg'd. Mxp'd. rami. Exopod of 2nd maxillae, its basipodite made up longer than 2nd maxillae, its basipodite made up of 2 elongated segts; end segt with 3 elongated but not claw-like setae. legs 1-seg'd (2-seg'd in Rami. of 1st 4 pairs of legs biramose, 1-seg'd in ♀ the figures). 5th legs biramose, 1-seg'd in ♂ unknown. Length 5-7mm. Color whitish and transparent.

PSEUDOLOVENULA MARUKAWA, 1921 CARD 2
- MAUCHLINE 1988: 712

Diaptomidae - Calanoida

Pseudolorenula magna Marukawa, ♀.

— — Marukawa, 1921, p. 13 : pls. 1-2.

--, TAJSKA, 1956a : 264

Habitat. Northern Japan.