

Ascidicofidae - Notodelphyoidea.

Scolecimorpha Notodelphyidae: cyclopoida <sup>(M)</sup>

G. O. Sars, 1926

S. insignis.

Syn. Scolecimorpha Sars, 1926, p. 3.

- Illg + Dudley 1961: 87-91

Length 13 mm. !! . Body vermiform-cylindrical.  
Curved dorsally into a semicircle or arc. Head  
well-defined: rostrum obtuse: two large lateral fleshy  
expansions, rounded at posterior corners. A.  
rounded protuberance on either side just behind  
the head - segmentation of entire body obsolete -  
1st Ant. indistinctly 3-segd. 2nd Ant 2-segd, uni-  
ramose, mouth parts rudimentary - 4 pairs of  
biramose legs, very rudimentary, rami 1-segd -  
No external ovisacs: entire body cavity serving as  
a brood chamber,

Scolecimorpha huntsmani Henderson - see Scolecodes

Scolecimorpha insignis Sars 1926

-- Stock 1967e: 56-57, possibly a synonym of Ophioseides joubini. Chatton, 1909?

Scolecimorpha joubini (Chatton 1909)

- ---, Bresciani et Lutzen 1962. PP. 369-370, fig. 1  
--- ---, Illg & Dudley, 1961. pp. 87-91, fig. 29-30  
--- ---, Illg & Dudley, 1965, pp. 440-441.

- - DUDLEY + ILLG 1991: 6 -

Scolecimorpha joubini (Chatton)

--Jones J.B., 1973, p. 256

— Monniot, 1961: 97-100, fig. 1