Academy of Natural Sciences

A Parasitic Copepod Author(s): Prof. Leidy

Source: Proceedings of the Academy of Natural Sciences of Philadelphia, Vol. 41 (1889), p. 95

Published by: Academy of Natural Sciences Stable URL: http://www.jstor.org/stable/4061586

Accessed: 02/12/2009 01:34

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at http://www.jstor.org/page/info/about/policies/terms.jsp. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at http://www.jstor.org/action/showPublisher?publisherCode=ans.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact support@jstor.org.



Academy of Natural Sciences is collaborating with JSTOR to digitize, preserve and extend access to Proceedings of the Academy of Natural Sciences of Philadelphia.

APRIL 16.

The President, Dr. JOSEPH LEIDY, in the chair.

Thirty-four persons present.

The following papers were presented for publication:—

- "A contribution to the Spider fauna of the Bermuda Islands," by Dr. Geo. Marx.
- "Notes on a small collection of Myriapods from the Bermuda Islands," by Charles H. Bollman.
- "A contribution to the Actinology of the Bermudas," by J. Playfair McMurrich, Ph. D.
- "On some new species of Mollusca from the Bermuda Islands," by Angelo Heilprin.
- A Parasitic Copepod.—Prof. LEIDY remarked that last summer while at Beach Haven, N. J., there was brought to him from the surf a living specimen of the singular, transparent fish *Leptocephalus*. In examining it he observed attached to the tail-fin a minute cope-pod crustacean, apparently of the genus *Chalimus*. The parasite was attached by a long filiform rostrum and resembled in this and other respects more the Chalimus scomberi, as represented by Baird, in fig. 5, tab. xxxiii, of the British Entomostraca, than it does the



original of this species as represented by Burmeister in the Nova Acta N. C. of Bonn, xvii, tab. 23, fig. 13. The species which may be distinguished as *Chalimus tenuis* is considerably less than half the size of *C. scomberi*. The cephalothorax nearly twice the length of the breadth, is obcordate and proportionately much narrower than in the latter species. The frontal segment is narrow and not prominent laterally, and the biarticulate antennae are concealed beneath. The abdomen, half the length of the cephalothorax, exhibits three conspicuous divisions, and the short caudal appendages end in three minute setae. Abdominal feet ending in biramous leaf-like segments fringed with short setae. Rostrum linear and almost as long as the cephalothorax. Whole length 1.125 mm; length of cephalothorax 0.5 mm; breadth 0.275; length of rostrum 10.5; length of abdomen 0.25 mm.

> The accompanying outline represents the animal magnified forty-four diameters.